

**Table (2):** Clinical findings of studied groups.

Variable	Group (Ia) N = 25	Group (Ib) N = 25	Group (II) N = 25	X <sup>2</sup>	p- value
	N (%)	N (%)	N (%)		
<b>Presenting Symptoms</b>					
<b>Fever:</b>					
- Yes	18 (72.0%)	6 (24.0%)	14 (56.0%)	11.9	0.003 <sup>a, c</sup>
- No	7 (28.0%)	19 (76.0%)	11 (44.0%)		
<b>GI Bleeding:</b>				7.78	0.005 <sup>b, c</sup>
- Yes	9 (36.0%)	6 (16.0%)	0 (0.0%)		
- No	16 (64.0%)	19 (84.0%)	25 (100.0%)		
<b>Abdominal pain:</b>				3.95	0.14 NS
- Yes	7 (28.0%)	3 (12.0%)	9 (36.0%)		
- No	18 (72.0%)	22 (88.0%)	16 (64.0%)		
<b>Presenting Signs</b>					
<b>Jaundice:</b>				7.01	0.03 <sup>b</sup>
- Yes	10 (40.0%)	6 (16.0%)	2 (8.0%)		
- No	15 (60.0%)	19 (84.0%)	23 (92.0%)		
<b>Abdominal Tenderness:</b>				5.44	0.65 NS
- Yes	6 (24.0%)	1 (4.0%)	7 (28.0%)		
- No	19 (76.0%)	24 (96.0%)	18 (72.0%)		

**a:** group (Ia) vs group (Ib);

**b:** group (Ia) vs group (II);

**c:** group (Ib) vs group (II)