

**Table (5):** Comparison between reactive and non-reactive anti-*Toxoplasma* IgG in patients with HD regarding residency, vascular access, lymphadenopathy, organomegaly, animal exposure, Age, the duration of dialysis, and CBC with the differential leukocytic count.

		Patients with non-reactive anti- <i>Toxoplasma</i> IgG (n=40)		Patients with reactive anti- <i>Toxoplasma</i> IgG (n=10)		Test of sig.		
		N	%	N	%	p-value	Sig.	
<b>Residency</b>	Urban	35	87.5%	9	90.0%	>0.999	NS	
	Rural	5	12.5%	1	10.0%			
<b>Vascular access</b>	Fistula	37	92.5%	9	90.0%	>0.999	NS	
	Others	3	7.5%	1	10.0%			
<b>Lymph-adenopathy</b>	No	34	85.0%	8	80.0%	0.653	NS	
	Yes	6	15.0%	2	20.0%			
<b>Organomegaly</b>	No	32	80.0%	4	40.0%	0.020	S	
	Yes	8	20.0%	6	60.0%			
<b>Exposure to animals</b>	No	32	80.0%	7	70.0%	0.671	NS	
	Yes	8	20.0%	3	30.0%			
		<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>p-value</b>	<b>sig.</b>	
<b>Age</b>		14.46	4.51	13.95	4.35	0.751	NS	
<b>The duration of dialysis</b>		7.19	5.10	7.65	4.75	0.798	NS	
		<b>Patients with non-reactive anti-<i>Toxoplasma</i> IgG (n=40)</b>		<b>Patients with reactive anti-<i>Toxoplasma</i> IgG (n=10)</b>		<b>t-test</b>		
<b>CBC with differential leucocytic count</b>		<b>Mean</b>	<b>SD</b>	<b>Mean</b>	<b>SD</b>	<b>value</b>	<b>p-value</b>	<b>sig.</b>
<b>Hemoglobin (13- 17) g/dl</b>		10.65	1.72	10.39	2.08	t = 0.41	0.687	NS
<b>Platelets (200-500x10<sup>3</sup>)/UL</b>		199.87	77.19	194.40	106.76	t = 0.18	0.854	NS
<b>TLC (5-15x10<sup>3</sup>)/UL</b>		6.55	2.13	6.89	1.91	t = -0.46	0.645	NS
<b>Neutrophils (2-8x10<sup>3</sup>)/UL</b>		3.66	1.62	3.94	1.67	t = -0.49	0.629	NS
<b>Lymphocytes (1-5x10<sup>3</sup>)/UL</b>		2.09	.79	2.09	1.04	t = 0.01	0.989	NS
		<b>Median</b>	<b>IQR</b>	<b>Median</b>	<b>IQR</b>	<b>Mann-Whitney test</b>		
<b>Eosinophils (0.2-1x10<sup>3</sup>)/UL</b>		0.25	0.1 - 0.58	0.37	0.2 - 0.5	U = 172	0.495	NS